Student Medical Exemption Certificate for Required Immunizations

The information below must be entered in the Department of Public Health Medical Exemptions portal. More information is available here.

Name of Primary Care Provider granting exemption: _____

Please check one (practitioner granting exemption must be licensed as one of the following):

☐ Physician (MD or DO) ☐ Physician Assistant ☐ APRN

CT License number: _____

NPI: _____

Phone number: _____ Email: _____

Directions:

Part 1. Please complete the demographics section on the patient/student.

Part 2. Please mark the contraindications/precautions that apply to this patient/student (indicate all that apply).

Part 3. If no contraindications or precautions apply in part 2, write a brief explanation of the reason the patient/student requires the exemption.

Part 4. Sign the Statement of Clinical Opinion and date the form.

Attach a copy of the patient/student's most current immunization record.

Part 1. Patient/Student Information:

First name (in full) _____ Middle initial _____ Last name _____

Date of Birth _____

Mailing Address _____ City _____

State _____ Zip _____

Parent/Guardian: First Name _____ Last name _____

Primary phone number _____

School name _____

School address

_____

City _____

State _____ Zip _____

Current or Grade student is entering _____

Part 2. Please mark the vaccine(s), exemption duration, and all contraindications/precautions that apply to this patient/student for each vaccine.

Medical contraindications and precautions for immunizations are based upon the Advisory Committee on Immunization Practices (ACIP) Comprehensive General Recommendations and Guidelines, published by the Centers for Disease Control and Prevention.

A **contraindication** is a condition in a recipient that increases the risk for a serious vaccine adverse event (VAE) or compromises the ability of the vaccine to produce immunity.

A **precaution** is a condition in a recipient that might increase the risk for a serious VAE or that might compromise the ability of the vaccine to produce immunity. Under normal conditions, vaccinations are deferred when a precaution is self-limiting, but can be administered if the precaution condition improves.

CDC Recognized Contraindications and Precautions

Vaccine	Exemption Duration	ACIP Contraindications and Precautions (Check all that apply)
☐ **Diphtheria-Tetanus-and acellular Pertussis (DTaP)**	☐ **Temporary through:** ____/_____ **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component ☐ Encephalopathy within seven days after receipt of previous dose of DTP or DTaP **Precautions** ☐ Progressive neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy: defer DTaP until neurologic status clarified and stabilized ☐ GBS <6 weeks after previous dose of tetanus-toxoid–containing vaccine ☐ Fever greater than 40.5°C (104.9°F) <48 hours after vaccination of previous dose of DTP or DTaP ☐ History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid–containing or tetanus-toxoid–containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid–containing vaccine ☐ Moderate or acute illness with or without fever
☐ **Hepatitis A**	☐ **Temporary through:** ____/_____ **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component **Precautions** ☐ Moderate or severe acute illness with or without fever

☐ **Hepatitis B**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component ☐ Hypersensitivity to yeast **Precautions** ☐ Moderate or severe acute illness with or without fever
☐***Haemophilus influenza*e type b (HiB)**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component ☐ Age <6 weeks **Precautions** ☐ Moderate or severe acute illness with or without fever
☐ **Inactivated Influenza Virus (IIV)**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after previous dose of influenza vaccine or to vaccine component **Precautions** ☐ GBS <6 weeks after a previous dose of influenza vaccine ☐ Moderate or severe acute illness with or without fever ☐ Egg allergy other than hives, e.g., angioedema, respiratory distress, lightheadedness, recurrent emesis; or required epinephrine or another emergency medical intervention (IIV may be administered in an inpatient or outpatient medical setting and under the supervision of a health care provider who is able to recognize and manage severe allergic conditions).
☐ **Inactivated Polio Vaccine (IPV)**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component **Precautions** ☐ Pregnancy ☐ Moderate or acute illness with or without fever

☐ **Live Attenuated Influenza Virus (LAIV)**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component ☐ Concomitant use of aspirin or aspirin-containing medication in children and adolescents ☐ LAIV4 should not be administered to persons who have taken oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days.(e) ☐ Pregnancy ☐ Children aged 2 through 4 years who have received a diagnosis of asthma or whose parents or caregivers report that a health care provider has told them during the preceding 12 months that their child had wheezing or asthma or whose medical record indicates a wheezing episode has occurred during the preceding 12 months. ☐ Persons with active cerebrospinal fluid/oropharyngeal communications/leaks. ☐ Close contacts and caregivers of severely immunosuppressed persons who require a protected environment. ☐ Persons with cochlear implants (due to the potential for CSF leak, which might exist for some period of time after implantation. Providers might consider consultation with a specialist concerning risk of persistent CSF leak if an age-appropriate inactivated or recombinant vaccine cannot be used). ☐ Altered Immunocompetence ☐ Anatomic or functional asplenia (e.g. sickle cell disease) **Precautions** ☐ GBS <6 weeks after a previous dose of influenza vaccine ☐ Asthma in persons aged 5 years old or older ☐ Medical conditions which might predispose to higher risk of complications attributable to influenza(d) ☐ Moderate or severe acute illness with or without fever
☐ **Meningo-coccal conjugate vaccines (MenACWY)**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component, including yeast **Precautions** ☐ Moderate or severe acute illness with or without fever

☐ **Measles-Mumps-Rubella (MMR)**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component ☐ Pregnancy ☐ Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised) ☐ Family history of altered immunocompetence (i) **Precautions** ☐ Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product) ☐ History of thrombocytopenia or thrombocytopenic purpura ☐ Need for tuberculin skin testing or interferon-gamma release assay (IGRA) testing (k) ☐ Moderate or severe acute illness with or without fever
☐ **Pneumo-coccal (PCV13)**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose of PCV13 or any diphtheria-toxoid–containing vaccine or to a component of a vaccine (PCV13 or any diphtheria-toxoid–containing vaccine), including yeast **Precautions** ☐ Moderate or acute illness with or without fever
☐ **Tdap**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component ☐ Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, or Tdap **Precautions** ☐ GBS <6 weeks after a previous dose of tetanus-toxoid–containing vaccine ☐ Progressive or unstable neurological disorder, uncontrolled seizures, or progressive encephalopathy until a treatment regimen has been established and the condition has stabilized ☐ History of Arthus-type hypersensitivity reactions after a previous

		dose of diphtheria-toxoid—containing or tetanus-toxoid–containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid–containing vaccine ☐ Moderate or severe acute illness with or without fever
☐ **Varicella**	☐ **Temporary through:** **____/_____** **mm/ yyyy** ☐ **Permanent**	**Contraindications** ☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component ☐ Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised) (g) ☐ Pregnancy ☐ Family history of altered immunocompetence (j) **Precautions** ☐ Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product) ☐ Moderate or acute illness with or without fever

Part 3. Other Allergic Reactions/ Other Type of Medical Condition

Complete this section if claiming a medical exemption for a vaccine based on a condition that does NOT meet any of the ACIP criteria for a contraindication or precaution listed in part 2.

Vaccine(s), list all that apply:

For each vaccine listed above, select the allergic or other reaction for which medical exemption is being submitted. Please check off any of the following that apply:

☐ This patient has an autoimmune disorder

☐ This patient has a family history of an autoimmune disorder

☐ This patient has a family history of a reaction to a vaccination

☐ This patient has a genetic predisposition to a reaction to a vaccination as determined through genetic testing

☐ This patient has a previous documented reaction that is correlated to a vaccination

☐ Other condition/reaction not listed above (must specify): _____

Please provide an explanation of the reaction/condition listed above:

_____

Part 4. Statement of Clinical Opinion

In accord with the legal requirements of Public Act 21-6, the vaccine(s) indicated above is/are in my clinical opinion medically contraindicated for this patient/student due to the physical condition as explained above.

Clinician's Signature_____

Date _____

A person may be placed into quarantine or isolation when there are "reasonable grounds to believe [a person] to be infected with, or exposed to, a communicable disease or to be contaminated or exposed to contamination or at reasonable risk of having a communicable disease or being contaminated or passing such communicable disease or contamination to other persons if the commissioner determines that such individual or individuals pose a significant threat to the public health and that quarantine or isolation is necessary and the least restrictive alternative to protect or preserve the public health." Conn. Gen. Stat. § 19a-131b(a).